EXtrelut[®] NT

試料前処理用 液-液抽出カラム
エキストレルート NT

EXtrelut[®] NT

試料前処理用 液一液抽出カラム
エキストレルート NT

目次

ページ

● 抽出原理・構造	1
● 試料と溶媒	2
● 特徴と利点	2-4
● 操作法	4-5
● 具体的使用例	
・尿からの抽出	5-6
・血液成分からの抽出	7-8
・組織や排泄物からの抽出	8
● エキストレルート NT を用いた 実際の分析例	9
● 参考文献リスト	10-19
● エキストレルート NT 製品ラインアップ	20

試料前処理用 液-液抽出カラム エキストレルート NT

生体試料中の脂溶性物質の分析には、実際の定性・定量分析に先立って、サンプルの前処理が必要になります。一般に用いられている栓付き試験管もしくは分液ロートを用いた液-液抽出法では、振とうや遠心分離等の煩雑な操作が必要で、また大量の有機溶媒を消費することになります。さらにサンプルによってはエマルジョンが形成し、層分離が不完全であったり、目的物質の回収率が著しく低下することがあります。エキストレルートNTはワイドポア (巨大細孔) をもつ珪藻土 (キーゼルグール) を特別に調製し、充填剤として使用した試料前処理用液-液抽出ディスプレイカラムです。このカラムを用いることで上記のような問題点は解決され、試料中から簡単に効率のよい液-液抽出を行なうことが可能になりました。

抽出の原理

水系試料をエキストレルートNTにアプライしますと、分析対象物質を含んだ試料中の水分が化学的に不活性な担体 (珪藻土) の周りに分配され、薄いフィルムを形成し、固定相のように作用します。

続いて、水とは混合しない有機溶媒 (例えばジエチルエーテル、酢酸エチル、ハロゲン化炭化水素) を流しますと、珪藻土表面にできた水層中に存在する脂溶性物質がその有機溶媒によって抽出され、カラムから溶出されます (図1)。

構造

エキストレルートNT1およびNT3はガラスカラムに、エキストレルートNT20は高密度ポリエチレン (HDPE) 製カラムに珪藻土が詰まっております。その基本構造は図2の通りです。

試料と溶媒

従来の液-液抽出を行なうことができる試料はすべて、エキストレルートNTを用いた前処理が可能です。また、エキストレルートNTはpH1~13と広いpH範囲の試料にお使いいただけます。

例えば、

- 尿
- 全血、血漿、血清
- その他の生体溶液（胃液、髄液、羊水など）
- 動植物、微生物の抽出物
- 農工業用水、排水
- 乳・乳製品
- 飲料（ジュース類）
- 酒類（ビール、ワイン、酒）
- 野菜、果物類
- 医薬品（錠剤、軟膏、シロップ剤、生薬など）

基本的には液-液抽出に用いられる有機溶媒はすべて使用できます。（下記参照）

また、抽出性の良い混合溶媒*としては次のようなものが挙げられます。

- **エーテル類**
ジエチルエーテル、ジイソプロピルエーテル、
t-ブチルメチルエーテル
- **エステル類**
酢酸エチル、酢酸メチル
- **炭化水素類**
ヘキサン、シクロヘキサン、ベンゼン
- **ハロゲン化炭化水素類**
クロロホルム、ジクロロメタン、ジクロロエタン
- クロロホルム／メタノール (90:10~85:15)
- ジエチルエーテル／エタノール (90:10~80:20)
- 酢酸エチル／2-プロパノール (90:10~80:20)
- ジクロロメタン／2-プロパノール (90:10~85:15)

* 混合溶媒を用いる際に低級脂肪族アルコール（メタノール、エタノールなど）のような極性溶媒の比率を高めると、珪藻土に保持された水相の体積が増加し、カラムから水分が脱着してしまう可能性がありますのでご注意ください。

特徴と利点

処理時間を大幅に短縮できる

従来の分液漏斗などを用いた液-液抽出では、抽出のための振とうや液相分離のための遠心分離など、煩雑な操作が必要になるので、一度に多数の試料を処理することが困難です。エキストレルートNTはディスポーザブルタイプのカラムで、試料の処理に振とうや遠心分離などの操作が必要ありません。また抽出後の有機溶媒における脱水操作も省くことができます。従いまして、エキストレルートNTを使用することで、多数の試料を簡単に短時間で処理することが可能になります。表1に従来の液-液抽出法とエキストレルートNTを用いた方法での、処理操作ステップ数やサンプルの処理時間の比較を示します。

	従来の液-液抽出		エキストレルートNTによる抽出		
	2	10	2	10	30
操 作	サンプル調製 ↓ 振とう抽出 (2-3回) ↓ 二相分離 (放置、遠心分離) ↓ 脱 水 ↓ 濃 縮		サンプル調製 ↓ カラム内で抽出 (1回) ↓ 濃 縮		
処 理 時 間	1時間	5時間	20分	1時間15分	2時間15分
1 試料あたりの処理時間	1時間	30分	10分	7.5分	4.5分

表1 従来の液-液抽出法とエキストレルートNTを用いた方法との比較 その1

エマルジョンを形成しない

生体試料を有機溶媒で抽出する場合、従来の液-液抽出法ではしばしばエマルジョンを形成し、層分離が不完全になり、目的物質の回収率の著しい低下や水溶性の不要成分の混入が生じます。エキストレルートNTを用いると、このようなエマルジョン形成はまったく見られず、以下に示すような、高い回収率や高い抽出効率を得られ、また夾雑物の混入も低く抑えることができます。

高い回収率を維持しつつ、溶媒の節約が行なえる

従来の定量的な液-液抽出では通常、有機溶媒による抽出を2回以上行なうため、大量の有機溶媒が使用されます。またエマルジョンの形成や低抽出効率によって、回収率が低くなりがちです。エキストレルートNTを用いると、定量的抽出が1回のカラム操作で行なうことができ、使用溶媒量も少量で済みます。表2に従来の液-液抽出法とエキストレルートNTを用いた方法での回収率と使用溶媒量の比較を示します。

目的物質 (抽出溶媒)	方法	試料容量 (ml)	回収率 (%)		総回収率 (%)	溶媒使用量 (ml)
			1回目	2回目		
バルビタール (エーテル)	従来法	40	85	13	98	80
	エキストレルートNT	20	99	—	99	35
モルフィン (ジクロロメタン/ 2-プロパノール 85 : 15)	従来法	40	(合計100)		100	80
	エキストレルートNT	20	99	5	100	55
コデイン (ジクロロメタン/ 2-プロパノール 85 : 15)	従来法	40	(合計100)		100	80
	エキストレルートNT	20	98	1	99	55

表2 従来の液-液抽出法とエキストレルートNTを用いた方法との比較 その2

夾雑物の混入を抑えることができる

エキストレルートNTではエマルジョンの形成が見られませんので、それに由来する夾雑物の混入がなく、純度の高い抽出物が得られます。また同時に、相対的な検出感度の向上が見られます。図3に従来の液-液抽出法とエキストレルートNTを用いた方法で処理した後の試料をガスクロマトグラフィーで分析した結果を示します。また表3に両方法で処理された試料における検出感度の比較を示します。

図3 従来の液-液抽出法とエキストレルートNTを用いた方法との比較 その3

両方法を用いた血清からの抽出物中の塩基性物質 (Eu=オイパベリン、Pa=パパベリン) を、硫酸洗浄により分離後、ガスクロマトグラフィーによって分析。

分析条件

サンプル量 : 5 μ l

カラム : クロモソルブG AW-DMCS

キャリアガス : N₂ (30ml/min)

昇温 : 170-275°C (8°C/min)

検出 : FID

目的物質 (抽出溶媒)	従来法 (pH9.4)	エクストレルートNT (pH8.9)	検出試薬
モルフィン	0.5 μ g	0.3 μ g	硫酸/ヘキサヨード白金酸
コデイン	1 μ g	0.6 μ g	硫酸/ヘキサヨード白金酸
フェノバルビタール	1 μ g	0.5 μ g	ジフェニルカルバゾン/硫酸水銀

表3 従来の液-液抽出法とエクストレルートNT を用いた方法との比較 その4

広いpH使用範囲

エクストレルートNTは化学的に不活性な充填剤を用いているため、pH1~13の範囲で安定で、最適な pH条件で抽出できます。またカラム内の pH を順次変化させることにより、1つの試料から酸性・中性・塩基性の条件でそれぞれ抽出可能です（ページ6参照）。

操作法

① 一般的操作法 (エクストレルートNT20使用の場合)

図4にエクストレルートNT20を用いる時の一般的操作法を示します。エクストレルートNT1もしくはNT3を使用される場合の操作も基本的に同じとなりますが、処理可能な試料量等が異なってきます。詳しくは表4をご参照ください。

図4 エクストレルートNT カラムの操作法

● 操作上の注意 ●

- 抽出時間は試料のタイプ、pHおよび溶出有機溶媒の種類によって異なってきます。
- NT20の場合、40mlの溶出溶媒を用いると抽出液は25mlとなります(カラムのデッドボリュームは15mlです)。
- オーバーロードによるカラムからの試料の流出を防ぐために、指定の試料量はお守りください。
- 試料量が少なすぎる場合は、試料を適量に希釈し、最大処理量に合わせてからアプライしてください。
- 抽出前の放置時間が短すぎると回収率が悪化することがあります。

エクストレルートNTの種類	(推奨される) 注射針のサイズ (内径 x 長さ mm)	処理可能な最大試料量 (ml)	抽出前の放置時間 (分)	推奨される溶出有機溶媒量 (ml)
NT1	0.6 x 30	1	5~10	6
NT3	0.6 x 30	3	5~10	15
NT20	0.7 x 30	20	10~15	40

表4 エクストレルートNT の種類と試料量

② 有機溶媒から塩形成物質を除去する時の操作法

③ 有機溶媒で抽出後、カラム内の残存水溶性試料を抽出する時の操作法

具体的使用例 ～NT20を用いて～

① 尿からの抽出

1. 酸性での抽出（酸性物質ならびに中性物質の抽出）

尿20mlを塩酸でpH1.5～2.0に調整し、カラムにアプライします。試料が十分浸透するまで（10～15分間）放置し、適当な有機溶媒40mlで抽出します。使用有機溶媒としてはジメチルエーテル、クロロホルム、ジクロロメタンなどが適しています。

2. 弱塩基性での抽出（塩基性物質の抽出）

尿20mlを塩化アンモニウム／アンモニアバッファー(pH9.5)* 1mlでpH8.5～9.5に調整し、カラムにアプライします。試料が十分浸透するまで（10～15分間）放置し、適当な有機溶媒40mlで抽出します。使用有機溶媒としてはクロロホルム、酢酸エチル、ジクロロメタン／イソプロパノール（85：15）などが適しています。

* 塩化アンモニウムの冷飽和水溶液を25%アンモニウム溶液でpH9.5に調整します。

限られた数の薬剤あるいは鎮痛剤のスクリーニングでは、弱塩基性での酸性、中性、塩基性物質の同時抽出が便利です。

図7 および表5 に各薬剤の抽出におけるpH依存性を示します。

図7 残存水溶液試料の抽出液

化合物	pH 2.0		pH 8.5~9.0	
	1回目の抽出	2回目の抽出	1回目の抽出	2回目の抽出
Morphine	—	—	80—95	5—20
Quinine	—20	60	90—100	—
Cocaine	60—70	20	80—100	—
Codeine	—	20	80—100	—
Methadone	80—90	—	80—100	—
Nicotine	—	—	80—100	—
Amitypytline	80—90	—	80—100	—
Papaverine	80—90	—	80—100	—
Diphenhydramine	80—100	—	80—100	—
Phenmetrazine	—	—	80—100	—
Mescaline	—	—	80	10
Norephedrine	—	—	80	20
Ephedrine	20	40	60	30
Amphetamine	—	—	80—90	—
Mephentermine	—	—	80—100	—
Oxazepam	80—100	—	80—100	—
Chlordiazepoxide	80	20	80—100	—
Nitrazepam	80—100	—	80—100	—
Diazepam	80—100	—	80—100	—
Methaqualone	80—100	—	80—100	—
4-Aminophenazone	—	—	80—100	—
Aminopyrine	40	40	80—100	—
Salicylic acid	80—100	—	—	—
Hexobarbital	80	20	80—90	—
Amobarbital	80	20	70—80	20
Thiopental	80—100	—	80—90	—
Phenobarbital	80	20	70—80	20
Secobarbital	80	20	70—80	20
Pentobarbital	80	20	70—80	20
Barbital	80—100	—	60—70	20
Chlorpromazine	60—70	20	80—90	10

表5 pH値に対する抽出率の依存度

3. 酸性及び弱塩基性での連続抽出

尿20mlに約0.5gの塩化アンモニウムを加え、25%塩酸でpH2に調整した後、カラムにアプライします。試料が十分浸透するまで(10~15分間) 放置し、ジエチルエーテル等の適当な有機溶媒40mlを流すことで、酸性ならびに中性物質を抽出します。さらにカラムに残存している溶媒を軽く減圧吸引することでそれらの物質を完全に回収します。

次に飽和アンモニアガスをカラム内に通し、カラム内のpHを酸性から塩基性に変化させた後、ジクロロメタン/イソプロパノール (85:15) 等の適当な有機溶媒40mlを流し、塩基性物質を抽出します。

4. 抱合体からの抽出

薬物や毒物は生体内で代謝され、代謝物やその抱合体として尿中に排泄されます。例えば、ベンゾジアゼピン、メタカロン、フェニチアジン、モルフィン、コデイン等は、グルクロン酸あるいは硫酸の抱合体として排泄されます。これらの抱合体から薬物を抽出するにはまず、抱合体のグルクロニダーゼ/アリルスルファターゼによる酵素的加水分解、もしくは、塩酸存在下での加熱加水分解が必要となります。

抱合体の酵素加水分解は、尿20mlに2M酢酸ナトリウム／酢酸バッファー

(pH5.5) 2mlとグルクロニダーゼ／アリルスルファターゼ*50 μ lを混合し、40～45 $^{\circ}$ Cで一晩(約16時間)インキュベートして行います。塩酸加水分解は、尿10mlに25%塩酸10mlを加え還流下で15分間加熱することで行い、冷却後徐々に水酸化ナトリウム2～3gを加えて中和します。

これらの加水分解で調製された試料20mlを塩化アンモニウム／アンモニアバッファー (pH9.5) 1～2mlでpH8.5～9.5に調整し、カラムにアプライします。試料が十分浸透するまで(10～15分間)放置し、ジクロロメタン／イソプロパノール(85：15) 40mlで抽出します。

上記の方法ですと、抱合体中の代謝薬物のみならず遊離型代謝薬物並びに未代謝薬物も合わせて抽出されます。抱合代謝薬物のみを抽出したい場合は図8に示すように、まず抱合体のみを分離し、それを同様な方法で加水分解した後に抽出を行います。

* グルクロニダーゼ／アリルスルファターゼ溶液が
ございます
(製品番号104114)

図8 抱合代謝薬物の抽出法

② 血液成分からの抽出

血液成分(血液、血清、血漿)は、高タンパク質含量によって粘度が高いため、エキストレルートNTカラムにアプライする前に試料の希釈が必要となります。表6に血液からの各薬物の抽出効率に及ぼす希釈率の影響を示します。また溶出に使用される溶媒には試料の凝固を起こさない溶媒が適しており、ヘプタン、ジエチルエーテル、ジイソプロピルエーテル、酢酸メチル、酢酸エチル、ジクロロメタン等が用いられます。

目的物質(薬物)*	希釈割合(試料+水)	pH	回収率(%)	溶出溶媒 (希釈血液20mlに対する溶媒量)
モルフィン	1 + 2	9	65	酢酸エチル (50ml)
	1 + 4		78	
	1 + 9		92	
パパベリン	1 + 2	8～11	94	酢酸エチル (50ml)
	1 + 4		97	
	1 + 9		93	
レボプロマジン	1 + 2	9	57	酢酸エチル (50ml)
	1 + 4		65	
	1 + 9		56	
ニトラゼバム	1 + 1	9.5	79	エーテル (50ml)
オキサゼバム	1 + 1		86	
クロルジアゼポシド	1 + 1		81	
バルビタール	1 + 4	3	95	エーテル (50ml)
フェノバルビタール	1 + 4	3	90	
フェノバルビタール	1 + 4	3	88	

* 薬物濃度は全血に対し600ng～200 μ g/ml

表6 エキストレルートNTによる新鮮な全血の抽出

1. 血清からの抽出

酸性物質の抽出には0.1M塩酸溶液を、塩基性物質の抽出には0.1M水酸化ナトリウム溶液を、試料と同量加え、希釈します。アルコール含有溶媒で抽出を行う場合は試料と同量の飽和食塩水で希釈します。希釈された試料20mlをカラムにアプライし、適当な溶媒40～60mlで抽出します。使用有機溶媒としてはジエチルエーテル、酢酸エチル、ジイソプロピルエーテル／酢酸メチル（70：30）、ジクロロメタンなどが適しています。

2. 血液、もしくは血漿からの抽出

血液(全血)は最低でも2倍に、通常は5倍に水で希釈します(この際、溶血が起きます)。凝固した血液は希釈前にホモジネートしておきます。酸性物質の抽出は希釈試料4mlに水15mlと1M塩酸溶液1mlを、塩基性物質の抽出には希釈試料4mlに0.025%アンモニア溶液16mlを加えた後、10～15分放置してからカラムにアプライします。粘度が高い時はカラムの出口側から軽く減圧吸引します。試料が浸透するまで(10分間)放置し、適当な有機溶媒40～60mlで抽出します。使用有機溶媒としてはジエチルエーテル、酢酸エチル、ジイソプロピルエーテル／酢酸メチル(70：30)などが適しています。

③ 組織や排泄物からの抽出

試料を硫酸アンモニウム、酒石酸、およびエタノールと混合し、ホモジネートします。これをろ過して、ろ液を乾固した後、水で溶解します。この試料溶液20mlを用いて、①で示したような方法で抽出を行います。

エキストレルートNTを用いた実際の分析例

ーヒト血清中の抗てんかん薬の分析ー

血清500 μ lをりん酸バッファー*500 μ lで希釈

エキストレルートNT1にアプライし、8分間放置

ジクロロメタン/2-プロパノール (9 : 1)
1mlを添加し、10分間放置

ジクロロメタン/2-プロパノール (9 : 1)
6mlを添加し、抽出

抽出物を窒素ガスにより乾固

乾固物をメタノール1mlで溶解

10 μ lをHPLCで分析

HPLC conditions :

Column : LiChroCart® 250-4
LiChrospher® RP-select B (5 μ m)

Mobile phase : A-Water / Acetonitrile (1 : 1)
B-Water

Gradient :	Time	%A	%B
	0 min	10%	90%
	30 min	60%	40%
	44 min	60%	40%
	44.1 min	100%	0%
	50 min	100%	0%
	51 min	10%	90%
	75 min	10%	90%

Flow : 1 mL / min

Temperature : 30°C

Detection : UV205nm

* りん酸2水素ナトリウム17.6g、りん酸水素2ナトリウム2水和物4.5g、アジ化ナトリウム1.5gを水1L中で溶解 (pH6.0~6.1)

サンプルならびに回収率

1. Ethosuximide	84 $\pm$ 7%
2. Primidone	100 $\pm$ 2%
3. α -methyl- α -propylsuccinimide	internal standard
4. Phenobarbital	96 $\pm$ 2%
5. Hexobarbital	99 $\pm$ 2%
6. Carbamazepine	97 $\pm$ 1%
7. Phenytoin	100 $\pm$ 1%

参考文献リスト

Compound	Title	Journal
Acetaminophen, Ethenzamide, Bromovalerylurea	EXTrelut NT column extraction for screening of drugs in biological materials by GC and GC-MS. A fatal poisoning by ingestion of a drug containing acetaminophen, ethenzamide and bromovalerylurea.	Nippon Hoigaku Zasshi, 1985, 39 (5), 386-391.
Acetophenone, Salicylic acid, Toluidine etc	Application of liquid-liquid partition chromatography to a student exercise in organic chemistry. III. Separation of mixtures using an EXTrelut NT column.	Kagaku to Kyoiku, 1986, 34 (3), 237-240.
N-Acetylglutamate	N-Acetylglutamate synthetase: Enzyme assay in human liver.	J. Clin. Chem. Clin. Biochem., 1982, 20 (5), 329-329.
6-Acetylmorphine	Determination of 6-acetylmorphine in urine as a specific marker for heroin abuse by HPLC with fluorescence detection.	Anal. Chim. Acta, 1985, 170 (1), 13-20.
6-Acetylmorphine, Morphine	Detection of 6-acetylmorphine in vitreous humor and cerebrospinal fluid : comparison with urinary analysis for proving heroin administration in opiate fatalities.	J. Anal. Toxicol., 1999, 23(3), 168-172.
Adriamycin	Determination of adriamycin and its fluorescent metabolites in biological fluids in in-patients with lung cancer by HPLC.	Chem. Pharm. Bull., 1985, 33 (9), 3868-3874.
Aflatoxin M ₁	Procedure for TLC determination of aflatoxin M ₁ in milk in the parts per 10 ¹² range.	J. Chromatogr., 1979, 178 (2), 543-549.
Alanycarb	Determination of alanycarb in agricultural products by HPLC.	Shokuhin Eiseigaku Zasshi, 1998, 39 (3), 218-224.
Alkaloids	Photometric micro-determination of the total alkaloid content in cytosine-containing alkaloid mixtures in plants by an improved Reifer's reagent.	Fresenius'Z. Anal. Chem., 1984, 319 (4), 434-435.
Aloin (Aloe emodin)	Metabolism of aloin - the influence of nutrition.	J. Pharm. Biomed. Anal., 1996, 14 (8-10), 1335-1338.
Alprazolam	HPLC method for the determination of alprazolam in plasma using the column-switching technique.	J. Chromatogr. Biomed. Appl., 1992, 116 (J. Chromatogr., 578), 259-263.
Alprenolol, Metoprolol	Determination of alprenolol and metoprolol in plasma by column LC.	J. Chromatogr. Biomed. Appl., 1988, 78 (J. Chromatogr., 434), 265-270.
Amineptine	Determination of amineptine and its main metabolite in plasma by HPLC after solid-phase extraction.	J. Chromatogr. Biomed. Appl., 1990, 97, (J. Chromatogr., 532), 351-361.
Amines, carcinogenic	Quantitation of mutagenic/carcinogenic heterocyclic aromatic amines in food products.	J. Chromatogr., 1992, 592, 271-278.
Amines, heterocyclic aromatic	Simple methods for quantifying mutagenic heterocyclic aromatic amines in food products.	Carcinogenesis, 1990, 11 (9), 1597-1603.
Amines, mutagenic	Isolation by solid-phase extraction and LC determination of mutagenic amines in beef extracts.	J. Chromatogr. A, 1996, 719 (1), 203-212.
Amines, mutagenic heterocyclic	An efficient and convenient method for the purification of mutagenic heterocyclic amines in heated meat products.	Carcinogenesis, 1989, 10 (7), 1175-1182.
Aminofenitrothion, Aminofenitroxon	Detection of S-methylenitrothion, aminofenitrothion, aminofenitroxon and acetylaminothion in the urine of a fenitrothion intoxication case.	Forensic Sci. Int., 1989, 41 (3), 245-254.
9-Aminophenanthrene-derivatized free fatty acids	Fluorimetric HPLC of 9-aminophenanthrene-derivatized free fatty acids.	J. Chromatogr. Biomed. Appl., 1984, 30 (J. Chromatogr., 305), 261-270.
Amitraz	Rapid and sensitive quantitation of amitraz in plasma by GC with nitrogen - phosphorus detection and its application for pharmacokinetics.	J. Anal. Toxicol., 1991, 15 (3), 116-118.
Amphetamine sulfate	Detection of sympathomimetic central nervous stimulants with special reference to doping III. Column extraction with EXTrelut NT.	J. Chromatogr., 1978, 161, 360-365.
Anabolic steroid (19-nor-testosterone)	Determination of anabolic steroid 19-nor-testosterone [nandrolone] in bovine serum by GC-SIM-MS.	J. Chromatogr. Sci., 1993, 31 (10), 429-432.
Androstadienone, Testosterone	Simultaneous determination of testosterone and androstadienone (sex attractant) in human plasma by GC-MS with high-resolution selected-ion monitoring.	J. Chromatogr., 1991, 565 (1-2), 35-44.
Androstene (4-Hydroxy-4-androstene-3,17-dione)	Quantitative determination of 4-hydroxy-4-androstene-3,17-dione, a potent aromatase inhibitor, in human plasma using isotope dilution MS.	J. Steroid. Biochem., 1989, 32 (5), 699-702.
Angiotensin converting enzyme inhibitor	Determination of a new angiotensin converting enzyme inhibitor and its active metabolite in plasma and urine by GC-MS.	J. Chromatogr. Biomed. Appl., 1987, 63 (J. Chromatogr., 419), 123-133.
Antagonist Z-300	Determination of new hydrogen receptor antagonist Z-300 in plasma by HPLC with fluorescence detection.	Yakugaku Zasshi, 1993, 113 (8), 586-590.
Anthocyanins	HPLC-MS application to anthocyanins of Vitis vinifera L.	J. Agric. Food Chem., 1995, 43 (8), 2104-2109.
Anthranilic acid	Micro-determination of anthranilic acid derivatives of anti-inflammatory drugs by HPLC and its application to forensic chemistry.	Eisei Kagaku, 1991, 37 (6), 461-466.
Anti-epileptic drugs	Mobile phase optimization in the separation of antiepileptic drugs by HPLC.	Ann. Clin. Biochem., 1986, 23 (5), 559-565.
Anti-epileptic drugs	Simultaneous determination of antiepileptic drugs and their metabolites, including chiral compounds, via beta cyclodextrin inclusion complexes by a column-switching-chromatographic technique.	J. Chromatogr. Biomed. Appl., 1994, 658 (2), 385-390.
Anti-epileptic drugs	An improved method for preparation of samples for the simultaneous assay of some antiepileptic drugs by GLC.	Clin. Chim. Acta, 1978, 90 (3), 203-208.
Anti-epileptic drugs	Procedure for LC determination of some commonly used antiepileptic drugs in serum.	Clin. Chim. Acta, 1978, 89, 301-309.
Anti-epileptic drugs	Sensitive analysis of antiepileptic drugs in very small portions of human brain by microbore HPLC.	J. Liq. Chromatogr., 1987, 10 (8-9), 1847-1864.
Anti-herpes agent	Metabolism of anti-herpes agent 5-(2-chloroethyl)-2'-deoxyuridine in mice and rats.	Eur. J. Drug. Metab. Pharmacokin., 1991, 16 (2), 129-136.
Anti-hypertensive drug	Determination of the antihypertensive drug 1-[2-ethoxy-2-(3-pyridyl)ethyl]-4-(2-methoxyphenyl)-piperazine in rat and human plasma by HPLC and isotope-dilution MS.	J. Chromatogr. Biomed. Appl., 1989, 79 (J. Chromatogr., 487), 331-340.
Anti-mycotic agents, Bifonazole	GC-MS determination of two new antimycotic agents in rabbit plasma following topical administration. A preliminary comparison with bifonazole.	J. Chromatogr. Biomed. Appl., 1991, 567 (1), 105-112.
Aromatic heterocyclic amines	Determination of heterocyclic aromatic amines by capillary zone electrophoresis in a meat extract.	J. Chromatogr. A, 1997, 769(2), 293-299.
Aromatic retinoic acid	Determination of the new aromatic retinoic acid Ro 13-7410 in plasma by two-dimensional GC- negative-ion CI-MS with selected-ion monitoring.	Biomed. Environ. Mass Spectrom., 1989, 18 (7), 453-463.

Compound	Title	Journal
Azaperone	Analysis and identification of azaperone and its metabolites in human.	J. Anal. Toxicol., 1992, 16 (1), 33-35.
Basic drugs	Impact of urine matrix and isolation procedure on retention behaviour of basic drugs in TLC.	J. Pharm. Biomed. Anal., 1988, 6 (4), 415-420.
Beer flavour components	Rapid method for the extraction and analysis of beer flavour components.	J. Inst. Brew., 1987, 93 (2), 113-115.
Benalaxyl residues	GC determination of benalaxyl residues in different crops and water.	J. Assoc. Off. Anal. Chem., 1993, 76 (3), 650-656.
Benazepril	Determination of benazepril, a new angiotensin-converting enzyme inhibitor, and its active metabolite, benazeprilat, in plasma and urine by capillary GC-MS detection.	J. Chromatogr. Biomed. Appl., 1988, 78 (J. Chromatogr., 434), 239-246.
Benzimidazole fungicides	Simultaneous determination of benzimidazole fungicides by HPLC on apples, pears and their pulps.	Pestic. Sci., 1989, 25 (4), 355-360.
Benzimidazolic fungicides	Simplified clean up for the determination of benzimidazolic fungicides by HPLC with UV detection.	J. Chromatogr. A, 1995, 697 (1-2), 145-152.
Benzodiazepines	Possibilities for the determination of some benzo-diazepine derivatives and a thienodiazepine by polarographic methods (differential pulse polarography and linear-sweep chronoamperometry) combined by an EXtrelut NT-column extraction in plasma or serum.	Sci. Pharm., 1984, 52 (1), 66-72.
Benzodiazepines	Quantitative LC, thermospray-tandem MS (LC - TSP - MS - MS) analysis of some thermolabile benzodiazepines in whole blood.	Forensic Sci. Int., 1992, 54 (1), 67-74.
Benzodiazepines	Determination of 2,3-benzodiazepine derivatives in rat plasma by HPLC.	J. Chromatogr. A, 1999, 846(1-2), 165-168.
Benzoic acid, Sorbic acid	Procedure for LC determination of food preservatives and Saccharin in foodstuffs.	J. Chromatogr., 1979, 173, 343-348.
Benzyl alcohol	Determination of benzyl alcohol and its metabolite in plasma by reversed-phase HPLC.	J. Chromatogr. Biomed. Appl., 1991, 106 (J. Chromatogr., 568), 145-155.
Benzylamine, Methyl salicylate	Application of liquid-liquid partition chromatography to the student exercise in organic chemistry. I. The extraction using an EXtrelut NT column.	Kagaku to Kyoiku, 1985, 33 (2), 152-155.
Benzylamine, Methyl salicylate	Application of liquid-liquid partition chromatography to the student exercise in organic chemistry. II. Comparison of methods for liquid-liquid extraction.	Kagaku to Kyoiku, 1985, 33, 344-345.
Beta agonists	Development and validation of a multi-residue method for beta agonists in biological samples and animal feed.	J. Assoc. Off. Anal. Chem., 1992, 75 (3), 554-560.
Beta blockers	Qualitative GC and GC-MS screening for beta blockers in urine after solid-phase extraction using EXtrelut NT columns.	J. Chromatogr. Biomed. Appl., 1988, 70 (J. Chromatogr., 426), 194-201.
Beta2-agonists, Cimeterol	Multiresidue analysis of beta2-agonists in human and calf urine using multimodal solid-phase extraction and HPLC with electrochemical detection.	J. Chromatogr. Biomed. Appl., 1999, 726(1-2), 149-156.
Beta-adrenergic agonist drugs	Synovial fluid as a matrix of selection in the detection of beta-adrenergic agonist drugs in carcasses and fresh meat.	Analyst, 1994, 119 (12), 2591-2593.
Beta-Blockers	Comparative study of extraction methods for the GC and GC-MS screening of urine for beta-blocker abuse.	J. Pharm. Biomed. Anal., 1987, 6 (6-8), 1988, 827-835.
Beta-blockers	Characterization of beta-blockers by chemical ionization witz ammonia: application to forensic chemistry.	Biol. Mass Spectrom., 1993, 22 (8), 462-464.
Biogenic amine-producing bacterial cultures	Rapid identification of biogenic amine-producing bacterial cultures using isocratic HPLC.	J. Chromatogr. A, 1996, 719 (2), 321-326.
Bopindolol	Column LC determination of hydrolysed bopindolol [4-(2-benzoyloxy-3-t-butylaminopropoxy)-2-methylindole hydrogen malonate], in the picogram per-millilitre range in plasma, using cartridge extraction and dual electrochemical detection.	J. Chromatogr. Biomed. Appl., 1987, 66 (J. Chromatogr., 422), 205-215.
Bromazepam	Determination of bromazepam in plasma and of its main metabolites in urine by reversed-phase HPLC.	J. Chromatogr. Biomed. Appl., 1984, 35 (J. Chromatogr., 310), 129-137.
Bromopropylate	GC determination of residues of bromopropylate and two of its degradation products in honey.	J. Assoc. Off. Anal. Chem., 1984, 67 (5), 896-898.
Bromvalerylurea	Separation and extraction of bromvalerylurea in body fluids by EXtrelut NT.	Hanzaigaku Zasshi, 1984, 50 (4), 153-156.
Bromvalerylurea	Rapid determination of bromvalerylurea in biological materials by HPLC.	Eisei Kagaku, 1986, 32 (5), 368-372.
Buflomedil	Quantitation and ultra-violet spectrum identification of buflomedil in whole blood and plasma by HPLC.	J. Anal. Toxicol., 1990, 14 (1), 18-21.
Buflomedil hydrochloride	GC-MS analysis of buflomedil hydrochloride in biological samples after acute intoxication.	Forensic Sci. Int., 1991, 51 (1), 125-129.
Buprenorphine	Analysis of buprenorphine in plasma and urine by GC.	J. Chromatogr. Biomed. Appl., 1990, 93 (J. Chromatogr., 528), 459-463.
Butyrophenones	Positive- and negative-ion MS of butyrophenones.	J. Chromatogr. Biomed. Appl., 1986, 55 (J. Chromatogr., 382), 135-145.
Cadralazine	HPLC determination of cadralazine in human plasma and urine.	J. Chromatogr. Biomed. Appl., 1988, 68 (J. Chromatogr., 424), 119-128.
Cafestol, Kahweol	Analysis of the content of the diterpenes cafestol and kahweol in coffee brews.	Food. Chem. Toxicol., 1997, 35(6), 547-554.
Caffeine	Simple and rapid method for the determination of caffeine in urine using EXtrelut NT columns.	J. Chromatogr. Biomed. Appl., 1983, 29, (J. Chromatogr., 278), 418-423.
Caffeine	Determination of caffeine in coffee, black tea and green tea by HPLC.	Shokuhin Eiseigaku Zasshi, 1988, 29 (2), 136-140.
Caffeine	Elimination of caffeine interference in HPLC determination of cotinine in human plasma.	J. Chromatogr. Biomed. Appl., 1991, 106 (J. Chromatogr., 568), 267-269.
Caffeine	Pharmacokinetics of caffeine and its metabolites in horses after intravenous intramuscular or oral administration.	Chem. Pharm. Bull., 1991, 39 (11), 2999-3002.
Caffeine, Theobromine	Simultaneous determination of purine alkaloids in daily foods by HPLC.	J. High. Resolut. Chromatogr., 1997, 20(8), 456-458.
Calcitriol	Scintillation proximity assay for calcitriol in serum without high-pressure LC.	Clin. Chim. Acta, 1993, 220 (1), 61-70.
Cannabinoids	Procedure for TLC determination of urinary cannabinoids.	J. Chromatogr., 1980, 196, 342-346.
Captopril	3-Bromomethylpropylphenazone as a new derivatization reagent HPLC of captopril and hydrochlorothiazide with UV detection.	Biomed. Chromatogr., 1998, 12(2), 57-60.

Compound	Title	Journal
Carbamate pesticides	Simple, rapid and simultaneous measurement of eight different types of carbamate pesticides in serum using LC - APCI-MS.	J. Chromatogr. Biomed. Appl., 1993, 131 (J. Chromatogr., 620), 61-71.
Carbamazepine	Chiral separation of 10,11-dihydro-10,11-trans-dihydroxycarbamazepine, a metabolite of carbamazepine with two asymmetric carbons in human serum.	J. Chromatogr. Biomed. Appl., 1996, 677 (2), 325-330.
N-4-Carbamoyl-3,4,5-dihydrodiazepam (uxepam)	Studies on the pharmacokinetics and metabolism of N-4-carbamoyl-3,4,5-dihydrodiazepam (Uxepam) in rats, dogs and man.	J. Pharm. Biomed. Anal., 1986, 4 (4), 497-503.
Carbaryl	Reversed-phase HPLC determination of carbaryl in post-mortem specimens.	J. Anal. Toxicol., 1985, 9 (4), 177-179.
Carbaryl, Propanil	HPLC determination of carbaryl and propanil contained in soft drinks.	Eisei Kagaku, 1983, 29 (4), 232-236.
Carbendazim, Thiabendazole	Selective clean-up applicable to aqueous acetone extracts for the determination of carbendazim and thiabendazole in fruits and vegetables by HPLC with UV detection.	J. Chromatogr. A, 1999, 833(1), 61-65.
Carboxylic acids	GC determination of organic acids from fruit juices by combined resin mediated methylation and extraction in supercritical carbon dioxide.	J. Chromatogr. A, 1997, 785(1-2), 251-261.
Cartap	Colorimetric and fluorometric determination of cartap by use of ion association with calcon.	Yakugaku Zasshi, 1983, 103 (6), 650-654.
Catechol estrogens	Assay of enzymic O-methylation of catechol estrogens by HPLC with coulometric detection.	J. Chromatogr. Biomed. Appl., 1993, 128 (J. Chromatogr., 617), 221-225.
Catechol, Guaiacol estrogens	Determination of catechol and guaiacol estrogens in urine by capillary GC-MS.	Biomed. Chromatogr., 1993, 7 (3), 172-176.
Catecholamines	Improves radioenzymic assay for the determination of catecholamines in plasma.	Clin. Chim. Acta, 1983, 132 (1), 111-116.
Cerebrovascular disorders agents (AJ 3941)	Determination of AJ 3941, a possible agent for the treatment of cerebrovascular disorders, in plasma and brain by means of HPLC with fluorescence detection.	J. Chromatogr. Biomed. Appl., 1992, 116 (J. Chromatogr., 578), 152-156.
Chlorambucil	Studies on the pharmacokinetics of chlorambucil and prednimustine in patients using a new HPLC assay.	Canc. Chem. Pharm., 1989, 23, 208.
Chloramphenicol	HPLC screening and confirmation methods for chloramphenicol residues in meat with off-line cartridge sample clean-up and on-line diode array UV-visible detection.	J. Chromatogr., 1986, 352, 445-453.
Chloramphenicol	Determination of chloramphenicol in tissues-problems with in vitro metabolism	Analyst, 1988, 113 (12), 1875-1876.
Chloramphenicol	Development and validation of a GC-MS procedure for the identification and quantification of residues of chloramphenicol.	Anal. Chim. Acta, 1990, 237 (1), 61-69.
Chlordane compounds and metabolite residues	Chlordane compounds and metabolite residues in termite control workers' blood.	Bull. Environ. Contam. Toxicol., 1995, 54 (6), 855-862.
Chlorhexidine	Selected-ion-monitoring assay for chlorhexidine in medical waste water.	Biomed. Mass Spectrom., 1984, 11 (5), 203-206.
Chlorophenoxyacetic acids	Simultaneous extraction and methylation of chlorophenoxyacetic acids from aqueous solution using supercritical carbon dioxide as a phase-transfer solvent.	Anal. Chem., 1994, 66 (24), 4459-4465.
Chloropropane diols	Sensitive method for the determination of 3-chloropropane-1,2-diol and 2-chloropropane-1,3-diol by capillary GC with MS detection.	J. Chromatogr. A, 1998, 802(2), 325-333.
Chloropropane diols	Determination of the chloropropanols, 3-chloro-1,2-propanediol and 2-chloro-1,3-propanediol, in hydrolysed vegetable proteins and seasonings by GC-ion trap tandem MS.	Rapid. Commun. Mass Spectrom. 1997, 11(13), 1417-1424.
3-Chloropropane-2-diol	Analysis of residual 3-chloro-2-propanediol in seasonings after derivatization with phenylboronic acid.	Shokuhin Eiseigaku Zasshi, 1995, 36 (3), 360-364.
3-Chloropropane-2-diol	Determination of 3-chloropropane-2-diol in hydrolysed vegetable proteins by capillary GC with electrolytic conductivity detection.	J. Chromatogr., 1993, 638 (1), 71-74.
Chloropropanols	Determination of chloropropanols in protein hydrolysates.	J. Chromatogr., 1992, 589 (1-2), 109-119.
Chloroquine	A rapid HPLC procedure for the simultaneous determination of chloroquine, monoethylchloro-quine, diazepam, and nordiazepam in blood.	J. Anal. Toxicol., 1989, 13 (2), 89-93.
Chlorpheniramine	Highly sensitive determination of chlorpheniramine as fluorescence derivative by HPLC.	J. Chromatogr. Biomed. Appl., 1987, 64 (J. Chromatogr., 420), 63-72.
Cholesterol	TLC of radioactively labelled cholesterol and precursors from biological material: a simple and sensitive method for investigating action on the sterol pathway.	Fresenius' Z. Anal. Chem., 1987, 327 (5-6), 558-560
Cimetidine	Rapid method for determination of cimetidine in biological fluids by HPLC using EXTrelut NT extraction.	J. Chromatogr. Biomed. Appl., 1983, 24 (J. Chromatogr., 273), 223-227.
Cisplatin	Analysis of mutagenic activity in human urine after concentration on different resins and HPLC.	Toxicol. Lett., 1989, 45 (2-3), 241-149.
Citaloprams	Simultaneous determination of citalopram, monodemethylcitalopram and didemethylcitalopram in plasma HPLC after column extraction.	J. Chromatogr. Biomed. Appl., 1990, 92, (J. Chromatogr., 527), 226-232.
Citrinin	HPLC determination of citrinin in cereals using an acid-buffered silica gel column.	J. Chromatogr., 1989, 462, 406-410.
Clenbuterol	Determination of clenbuterol in urine of calves by HPLC with in-series ultra-violet and electrochemical detection.	J. Pharm. Biomed. Anal., 1991, 9 (6), 485-492.
Clenbuterol, Salbutamol	Determination of clenbuterol and salbutamol in urine by capillary GC with capillary columns of 100 µm.	J. Chromatogr. A, 1993, 655 (1), 73-76.
Clomipramine	Improved clean-up procedure for the HPLC assay of clomipramine and its demethylated metabolite in human plasma.	J. Chromatogr. Biomed. Appl., 1995, 669 (2), 416-417.
Cloridine	Procedure for the GC determination of Clonidine in human plasma.	J. Chromatogr., 1980, 187, 161-169.
Cocaine, Benzoyl-ecgonine, local anaesthetic agents	LC analysis of cocaine, benzoyl-ecgonine, local anaesthetic agents and some of their metabolites in biological fluids.	J. Liq. Chromatogr., 1993, 16 (13), 2797-2811.
Cocaine, Ecgonine derivatives	Quantitation of cocaine, benzoyl-ecgonine and ecgonine methyl ester by GC-CI-MS-selected ion monitoring after EXTrelut NT extraction.	Forensic Sci. Int., 1984, 26 (3), 181-192.
Codeine, Morphine	Procedure for GC determination of Morphine and Codeine in urine.	Forensic Sci., 1976, 7, 131-140.
Corticosteroids	Chemiluminescence HPLC of corticosteroids using lucigenin as post-column reagent.	Biomed. Chromatogr., 1990, 4 (3), 119-122.
Cortisol	Procedure for radioimmunological determination of urinary free cortisol.	Clin. Chim. Acta, 1980, 101, 125-134.
Cortisol	Procedure for routinely practicable determination of free cortisol in urine by combination of automatic HPLC and RIA.	Clin. Chim. Acta, 1980, 106, 63-73.

Compound	Title	Journal
Cortisol	Pre-chromatographic derivatization of samples for (HP)TLC.	J. High Resolut. Chromatogr. Chromatogr. Commun., 1982, 5, 534-538.
Cotinine	Simultaneous determination of cotinine and trans-3'-hydroxycotinine in human serum by HPLC.	J. Chromatogr. Biomed. Appl., 1992, 115 (J. Chromatogr., 577), 358-361.
Cotinine	Urinary cotinine and exposure to parental smoking in a population of children with asthma.	Clin. Chem., 1999, 45(4), 505-509.
Cotinine, Nicotine	Hair analysis for nicotine and cotinine: evaluation of extraction procedures, hair treatments, and development of reference material.	Forensic Sci. Int., 1997, 84(1-3), 243-252.
Coumatetralyl	Screening for coumatetralyl in soft drinks by solid matrix extraction and HPLC with DAD.	J. Chromatogr., 1991, 553, 305.
p-Cresidine	Determination of p-cresidine [2-methoxy-5-methyl-aniline] in FD & C Red No. 40 by the diazotization and coupling procedure followed by reversed-phase HPLC.	J. Chromatogr., 1987, 405, 283-294.
Cresols	GC determination of cresols in the biological fluids of a non-fatal case of cresol intoxication.	Forensic Sci. Int., 1990, 47 (1), 21-29.
Cyclosporin A	Preparation, extraction and HPLC of cyclosporin A from plasma that contains interfering compounds.	Ann. Clin. Biochem., 1985, 22 (4), 402-407.
D&C Yellow No. 10	LC determination of 2-(2-quinoly) H-indene-3[2H]-dione and other organic-soluble matter in D&C Yellow No. 10.	J. Assoc. Off. Anal. Chem., 1985, 68 (3), 477-479.
Debrisoquine	LC method for the determination of debrisoquine and its 4-hydroxy-metabolite in human urine.	J. Chromatogr. Biomed. Appl., 1986, 51 (J. Chromatogr., 378), 509-513.
Demeton-S-methyl sulphoxide	Fast determination of demeton-S-methyl sulphoxide (Metasystox R) in blood plasma.	J. Anal. Toxicol., 1989, 13 (6), 330-332.
Deoxynivalenol	Oxidative transformation of deoxynivalenol (vomitoxin) for quantitative and chemical confirmatory purposes.	J. Agric. Food Chem., 1984, 32 (1), 72-75.
Deoxynivalenol	Analysis of trace levels of deoxynivalenol in cow's milk by high pressure LC.	J. Liq. Chromatogr., 1994, 17 (3), 673-683.
Dextromoramide	Determination of dextromoramide in plasma and whole blood using HPLC with ultra-violet absorbance detection.	J. Chromatogr. Biomed. Appl., 1992, 111 (J. Chromatogr., 573), 87-92.
Dextromoramide, Propoxyphenes	Simultaneous determination of dextromoramide, propoxyphene and norpropoxyphene in necropsic whole blood by LC.	J. Chromatogr. Biomed. Appl., 1993, 126 (J. Chromatogr., 615), 357-364.
Diacetoxyscirpenol	Simultaneous determination of diacetoxyscirpenol and T-2 toxin in fungal cultures and grain samples by capillary GC.	Food Addit. Contam., 1987, 4 (3), 277-284.
Dicyclomine hydro-chloride	Stability-indicating capillary GLC assay of dicyclomine hydrochloride in some pharmaceutical formulations.	J. Chromatogr., 1989, 475, 381-389.
Diethylene glycol	Determination of diethylene glycol in wine by use of an EXTrelut NT column.	Shokuhin Eiseigaku Zasshi, 1986, 27 (3), 283-287.
Diethylene glycol	Rapid determination of diethylene glycol in wine by GC using alumina column clean-up.	Shokuhin Eiseigaku Zasshi, 1986, 27 (2), 187-189.
Diethylene glycol	Determination of diethylene glycol in wine by high-performance liquid chromatography using anthracene-9-carbonyl chloride as a derivatizing reagent.	J. Chromatogr., 1988, 445 (2), 403-408.
Difenzoquat	Determination of difenzoquat in cereals by GC-MS.	Shokuhin Eiseigaku Zasshi, 1998, 39, 431-435.
Digoxin, Digoxigenin, Dihydrodigoxigenin	Determination of digoxin, digoxigenin and dihydrodigoxigenin in urine by extraction, derivatization HPLC.	J. Chromatogr. Biomed. Appl., 1986, 55 (J. Chromatogr., 382), 349-354.
Dihydroergotamine	Determination of sub-nanogram amounts of dihydroergotamine in plasma and urine using LC and fluorimetric detection with offline and online solid-phase drug enrichment.	J. Chromatogr. Biomed. Appl., 1987, 61 (J. Chromatogr., 417), 319-329.
Dihydroxyvitamin D ₃ , Vitamin D ₃	Development of a double-antibody radio-immunoassay for quantitation of 1 α ,25-dihydroxyvitamin D ₃ .	Clin. Chim. Acta, 1990, 189 (2), 97-110.
Dihydroxyvitamin D ₃ , Vitamin D ₃	Rapid, selective separation of 1 α ,25-dihydroxy-vitamin D ₃ from serum with EXTrelut NT columns.	Clin. Chem., 1987, 33 (1), 187.
Dihydroxyvitamin D ₃	Sensitive and simplified radio-immunoassay for 1,25-dihydroxyvitamin D ₃ .	Clin. Chim. Acta, 15. 1984, 142 (3), 325-338.
Dimethoate, Formothion	Determination of dimethoate in blood and hemo-perfusion cartridge following ingestion of formothion - case study.	Forensic Sci. Int., 1990, 48 (1), 79-88.
Disulfoton	Determination of disulfoton and its metabolites in the body fluids of a Di-Syston intoxication case.	Forensic Sci. Int., 1990, 48 (2), 145-154.
Diuretics	Detection of diuretics in horse urine by GC - MS.	J. Anal. Toxicol., 1992, 16 (3), 194-198.
Drugs	Procedure for GC determination of underivatized drugs in blood.	Clin. Chem., 1979, 25 (12), 2020-2025.
Drugs	Application of HPLC equipment with rapid-scan detection to the identification of drugs in toxicological analysis.	J. Liq. Chromatogr., 1994, 17 (19), 4131-4144.
Drugs	Identification of drugs of abuse by HPTLC.	J. Anal. Toxicol., 1980, 4, 141-144.
Drugs	Impact of biological matrix and isolation methods on detectability and interlaboratory variations of TLC Rf-values in systematic toxicological analysis.	J. Anal. Toxicol., 1984, 8 (4), 149-154.
Drugs	Rapid procedure for the isolation of drugs in biological samples: usefulness of enzymic digestion and the EXTrelut NT column.	Aichi Ika Daigaku Igakkai Zasshi, 15 (4), 1987, 617-625.
Dymron	Determination of dymron in agricultural products by HPLC.	Shokuhin Eiseigaku Zasshi, 1999, 40(3), 233-240.
Eicosapentaenoic acid	Fluorescence HPLC of eicosapentaenoic acid in serum and vital blood of fishes and in body fluid of plankton after labelling with 9-anthryldiazomethane.	Bunseki Kagaku, 1984, 33 (7), E271-E278.
Embutramide, Pentobarbitone	Determination of embutramide and pentobarbitone in meat and bone-meal by GC-MS.	Analyst, 1998, 123(12), 2513-2516.
Epomediol	Capillary GC determination of epomediol in human plasma and urine using a rapid solid-phase extraction.	Biomed. Chromatogr., 1989, 3 (5), 199-202.
Eproxindine hydrochloride	Determination of eproxindine hydrochloride in human plasma by capillary GC.	J. Chromatogr. Biomed. Appl., 1988, 71 (J. Chromatogr., 427), 157-161.
Estrogens	LC determination of the estrogens daidzein, formononetin, coumestrol, and equol in bovine blood plasma and urine.	J. Assoc. Off. Anal. Chem., 1988, 71 (5), 938-941.

Compound	Title	Journal
Ethopabate, Niicarbazine, Lasalocid sodium, Decoquinatone	Simultaneous determination of ethopabate, nicarbazine, lasalocid sodium and decoquinatone in mixed feeds by HPLC.	Shiryu Kenkyu Hokoku (Tokyo Hishiryu Kensasho), 1991, 16, 128-140.
Ethyl carbamate	Determination of ethyl carbamate in sake by alkylation and GC with a flame ionization detector.	Shokuhin Eiseigaku Zasshi, 1987, 28 (5), 330-335.
Ethyl carbamate	Simple analytical method for ethyl carbamate in alcoholic beverages.	Shokuhin Eiseigaku Zasshi, 1987, 28 (6), 492-497.
Ethyl carbamate	Measurement of ethyl carbamate in blood by capillary GC-MS using selected-ion monitoring.	Biomed. Environ. Mass Spectrom., 1990, 19 (1), 27-31.
Ethyl carbamate	Determination of ethyl carbamate in soy sauces.	Food Chem., 1993, 48 (3), 313-316.
Ethylamphetamine, stereoselective metabolism	Analysis of time-lapse changes of D- and L-enantiomers of racemic ethylamphetamine and stereoselective metabolism in rat urine by HPLC determination.	J. Anal. Toxicol., 1995, 19 (4), 225-228.
Ethylene thiourea	An improved method for residue analysis of ethylene thiourea in vegetables by HPLC.	J. Pestic. Sci., 1986, 11 (1), 81-84.
Ethylene thiourea, Propylene thiourea	Determination of ethylene thiourea and propylene thiourea in tomato products and in fruit purees.	Food Chem., 1993, 47 (3), 299-302.
Fatty acids	HPLC determination of fatty acids in phospholipid fraction of human platelets and plasma after administration of eicosapentaenoic acid.	Bunsei Kagaku, 1984, 33 (7), E219-E226.
Fatty acids, unsaturated	Procedure for TLC determination of radioactive labelled unsaturated compounds.	J. Chromatogr., 1979, 169, 213-222.
Fenitrothion	A rapid and sensitive method for detecting fenitrothion in biological fluids using the phosphorus-sulfur selective detector : a fenitrothion intoxication case.	Hiroshima J. Med. Sci., 1986, 35 (2), 87-92.
Flunarizine	Simple method for determination of flunarizine in serum by GC.	J. Chromatogr. Biomed. Appl., 1987, 65 (J. Chromatogr., 421), 372-376.
Fluoxetine, Norfluoxetine	Issues in methodology and applications for therapeutic drug monitoring of fluoxetine and norfluoxetine enantiomers.	Ther. Drug. Monit., 1998, 20(1), 20-24.
Food additives (t-Butylquinol)	Determination of food additives by HPLC. IV. Determination of t-butylquinol in oily foods and dried fish.	Shokuhin Eiseigaku Zasshi, 1984, 25 (2), 209-213.
Fungicides	Application of solid-phase partition cartridges in the determination of fungicide residues in vegetable samples.	J. Chromatogr., 1993, 643 (1-2), 363-368.
Fungicides	Rapid analysis of fungicides in agricultural products and processed foods by applying an accelerated solvent extraction system.	Shokuhin Eiseigaku Zasshi, 1997, 38(5), 358-362
Furazolidone	Determination furazolidone in swine plasma, muscle, liver, kidney, fat and urine based on HPLC separation after solid-phase extraction on EXTRELUT NT.	J. Chromatogr., 1986, 362 (1), 141-145.
General aspects	Determination of a new angiotensin converting enzyme inhibitor and its active metabolite in plasma and urine by GC-MS.	J. Chromatogr., 1987, 419, 122-133.
Ginsenosides	Procedure for LC determination of ginsenosides from ginseng drug preparations or plant material.	Planta Med., 1980, 39, 348-357.
Glyburide	Determination of glyburide in human plasma by LC with fluorescence detection.	J. Liq. Chromatogr., 1989, 12 (10), 1741-1758.
Glycerol trinitrate	Simultaneous determination of glycerol trinitrate and its two dinitrate metabolites in plasma and tissues by capillary GC	J. Chromatogr. Biomed. Appl., 1990, 99 (J. Chromatogr., 534), 196-201.
Granisetron	A rapid assay of granisetron in biological fluids from cancer patients.	Biol. Pharm. Bull., 1998, 21(5): 535-537.
Griseofulvin	Determination of griseofulvin in rat plasma by HPLC and HPTLC.	J. Chromatogr., 1979, 163, 322-326.
Growth-promoting compounds	Procedure for the GC-MS confirmation of some exogenous growth-promoting compounds in the urine of cattle.	J. Chromatogr., 1983, 279, 533-542.
Haloacetic acids	Measurement of haloacetic acids and their decomposition in wine by capillary GC.	Food Addit. Contam., 1991, 8 (1), 119-124.
Haloperidol, Hydroxyhaloperidol	HPLC analysis of haloperidol and hydroxyhaloperidol in plasma after solid-phase extraction.	J. Chromatogr. Biomed. Appl., 1990, 97 (J. Chromatogr., 532), 193-202.
Heterocyclic amines (HAA)	Determination of heterocyclic amines (HAA) in commercially available meat products and fish by HPLC-ESI-MS-MS.	Chromatographia, 1998, 48(3-4), 258-262.
Heterocyclic aromatic amines	Quantitative estimation of heterocyclic aromatic amines by ion-exchange chromatography and electrochemical detection.	J. Chromatogr. A, 1995, 697 (1-2), 377-382.
Heterocyclic aromatic amines	Evaluation of different clean-up procedures for the analysis of heterocyclic aromatic amines in a lyophilized meat extract.	J. Chromatogr. A, 1999, 836(2), 223-233.
Heterocyclic aromatic amines	Determination of heterocyclic aromatic amines in fried beefsteak, meat extract and fish by capillary zone electrophoresis.	Chromatographia, 1998, 48(9-10), 700-706.
Homopolymeric peptides	Fluorimetric determination of homopolymeric peptides with 5-dimethylaminonaphthalene-sulphonic acid after N-chlorination.	Chem. Pharm. Bull., 1989, 37 (10), 2841-2842.
Hydrocortisone	Clean up and re-use of kieselguhr (Extrelut) for liquid-liquid extraction of urinary cortisol.	J. Chromatogr. Biomed. Appl., 1997, 692(1), 238-240.
3-Hydroxybutyric acid	Determination of incubator-reject eggs in egg products by means of the 3-hydroxybutyric acid content. I. Improved enzymic and GC assays.	J. Agric. Food Chem., 1986, 34 (4), 663-666.
Hydroxycorticosteroids	Determination of 17-hydroxycorticosteroids in urine by fluorescence HPLC using Dns-hydrazine as a pre-column labelling reagent.	J. Chromatogr., 1982, 232 (1), 1-11.
4-Hydroxymandelic acid enantiomers	Determination of p-hydroxymandelic acid enantiomers in urine by HPLC with electrochemical detection.	J. Pharm. Biomed. Anal., 1997, 15(9-10): 1509-1514.
p-Hydroxymethadone	Synthesis, characterization and quantitation of para-hydroxymethadone, a metabolite of methadone in the rat.	Drug Metab. Dispos., 1986, 14 (4), 477-481.
4-Hydroxy-2-(4-methyl-phenyl)benzo-thiazole	Enzyme immunoassay for 4-hydroxy-2-(4-methylphenyl)benzothiazole.	J. Pharm. Biomed. Anal., 1993, 11 (6), 469-476.
4-Hydroxynonenal	Quantitative determination of the lipid peroxidation product 4-hydroxynonenal by HPLC.	Anal. Biochem., 1985, 150 (2), 369-378.
5-(4-Hydroxyphenyl)-5-phenylhydantoin	HPLC method for direct separation of 5-(4-hydroxyphenyl)-5-phenylhydantoin enantiomers using a chiral tris(4-methylbenzoate) column.	J. Chromatogr. Biomed. Appl., 1991, 106 (J. Chromatogr., 568), 157-163.

Compound	Title	Journal
Hydroxysteroids	Fluorescence HPLC determination of 3 alpha-hydroxysteroids in urine of 21-hydroxylase deficiency.	Biomed. Chromatogr., 1986, 1 (1), 1-6.
4-Hydroxytertatolol stereoisomers	Determination of 4-hydroxytertatolol stereoisomers in rat and human urine by HPLC.	J. Liq. Chromatogr., 1995, 18 (14), 2801-2812.
Hyoscyamine, Hyoscyne	Comparison between TLC-densitometry and HPLC for the determination of hyoscyamine and hyoscyne in leaves, fruit and seeds of <i>Datura</i> (<i>Datura</i> spp.).	J. Chromatogr., 1985, 329, 415-421.
Imipramine, Desipramine	Simultaneous determination of imipramine, desipramine and their deuterium labelled analogues in biological fluids by capillary GC-MS.	J. Chromatogr. Biomed. Appl., 1988, 70 (J. Chromatogr., 426), 93-101.
Iprnidazole, Ronidazole, Dimetridazole	LC multi-component methods for determination of residues of ipronidazole, ronidazole, and dimetridazole and some relevant metabolites in eggs, plasma, and faeces and its use in depletion studies in laying hens.	J. Assoc. Off. Anal. Chem., 1991, 74(1), 46-55.
Isosorbide dinitrate	Determination of isosorbide dinitrate and its mononitrate metabolites in human plasma using EXTrelut NT purification and capillary-column GLC.	J. Chromatogr. Biomed. Appl., 1984, 31 (J. Chromatogr., 306), 165-172.
Isosorbide nitrates	Clean-up of plasma extracts by GPC during analysis of isosorbide nitrates by capillary GC.	J. Chromatogr., 1991, 553 (1), 21-27.
Ketobemidone	Determination of ketobemidone in plasma by HPLC with electrochemical detection.	J. Chromatogr. Biomed. Appl., 1986, 48 (J. Chromatogr., 375), 207-211.
Lauric acid, Ethyl laurate	Simultaneous quantitation of lauric acid and ethyl laurate in <i>Sabal serrulata</i> by capillary GC and derivatization with trimethyl sulphoniumhydroxide.	J. Chromatogr. A, 1996, 719 (2), 479-482.
Laxatives	Qualitative and quantitative analysis of some synthetic, chemically acting laxatives in urine by GC-MS.	J. Chromatogr., 1981, 222 (3), 389-398.
Levamisole	Rapid EXTrelut NT column method for determination of levamisole in milk using HPLC.	J. Chromatogr. Biomed. Appl., 1985, 38 (J. Chromatogr., 337), 151-155.
Lidocaine	Toxicological analysis of lidocaine in biological materials by using HPLC: Benzocaine, procaine, tetracaine, bupivacaine.	Forensic Sci. Int., 1991, 49 (1), 65-73.
Local anaesthetics	Simultaneous determination of local anaesthetics including ester-type anaesthetics in human plasma and urine by GC-MS with solid-phase extraction.	J. Chromatogr. Biomed. Appl., 1999, 726(1-2), 185-194.
Lornoxicam and 5'-hydroxy-lornoxicam	Determination of the novel non-steroidal anti-inflammatory drug lornoxicam and its main metabolite in plasma and synovial fluid.	J. Chromatogr. Biomed. Appl., 1998, 707(1-2), 151-159.
Lysergide (LSD) and its N-demethyl metabolite	Determination of LSD and N-demethyl-LSD in urine by LC coupled to ESI-MS.	J. Chromatogr. Biomed. Appl., 1997, 692(2), 329-335.
Malathion	GC determination of malathion.	Eisei Kagaku, 1983, 29 (6), 423-426.
Methamphetamine	Separation of methamphetamine and amphetamine in blood and urine on an EXTrelut NT column.	Nippon Hoigaku Zasshi, 1983, 37 (1), 63-66.
Methamphetamine	Extraction of methamphetamine in human urine using EXTrelut NT column.	Kagaku Keisatsu Kenkyusho Hokoku, Hokagaku Hen. 37 (3), 1984, 224-226.
Methandienone	Detection of methandienone (methandrostenolone) and metabolites in horse urine by GC-MS.	J. Chromatogr. Biomed. Appl., 1992, 115 (J. Chromatogr., 577), 195-203.
Methomyl	Sensitive determination of methomyl in blood using GC-MS as its oxime tert-butylidimethylsilyl derivative.	J. Chromatogr. Biomed. Appl., 1998, 713(2), 323-330.
3-Methoxy-4-hydroxyphenylglycol	Determination of unconjugated 3-methoxy-4-hydroxyphenylglycol by LC for monitoring inhibition of monoamine oxidase activity in plasma.	Clin. Chem., 1987, 33 (11), 2078-2080.
3-Methoxy-4-sulfonyloxyphenylethanediol	Direct determination of 3-methoxy-4-sulfonyloxyphenylethanediol (MHPG sulphate) in urine using GLC.	J. Chromatogr. Biomed. Appl., 1984, 30 (J. Chromatogr., 305), 438-441.
Methylamphetamine	Simple extractive derivatization of methylamphetamine and its metabolites in biological materials with EXTrelut columns for their GC-MS determination.	J. Anal. Toxicol., 1997, 21(1), 54-58.
Methylamphetamine, Amphetamine	Estimation of optically active methylamphetamine and amphetamine using alpha-methoxy-alpha-trifluoromethylphenylacetyl chloride.	Nippon Hoigaku Zasshi, 1991, 45 (1), 1-5.
Methyl carbamate, Ethyl carbamate	Method for the determination of methyl carbamate and ethyl carbamate in wines.	Food Addit. Contam., 1992, 9 (2), 149-160.
N-Methyl carbamate pesticides	Determination of N-methylcarbamate pesticides in foods using an accelerated solvent extraction with a mini-column clean up.	Analyst, 1998, 123(4), 711-714.
Methylcitric acid	Methylcitric acid (2-hydroxybutane -2,3-tricarboxylic acid) determination in amniotic fluid by electron-impact mass fragmentography.	J. Clin. Chem. Clin. Biochem., 1988, 26 (5), 345-348.
Methylmalonic acid	Capillary GC-MS - stable isotope dilution analysis of methylmalonic acid in plasma: a candidate reference method for the diagnosis and the follow-up of methylmalonic acidaemias.	Chromatographia, 1993, 36, 212-214.
Methylparaben	Quantitative Gc and TLC determination of methylparaben (methyl hydroxybenzoate) in pharmaceutical dosage forms.	J. Chromatogr., 1984, 294, 431-435.
Methylprednisolone	Rapid method for the measurement of methylprednisolone and its hemisuccinate in plasma and urine following "pulse therapy" by HPLC.	J. Chromatogr. Biomed. Appl., 1985, 43 (J. Chromatogr., 342), 251-260.
Methylpyrazines	Simple and accurate determination of methylpyrazines in biofluids using HPLC.	J. Chromatogr., 1991, 567, 415.
Metoclopramide	Procedure for LC determination of metoclopramide in human serum.	Fresenius' Z. Anal. Chem., 1980, 301, 109-110.
Mexiletine	Investigation of blood level of mexiletine.	Byoin Yakugaku, 1987, 13 (3), 132-136.
Mexiletine	Sensitive determination method for mexiletine derivatized with dansyl chloride in rat plasma utilizing a HPLC peroxyoxalate chemiluminescence detection system.	J. Pharm. Biomed. Anal., 1997, 15(9-10): 1509-1514.
Microbial metabolites	Pre-column switching in HPLC quantitation of small amounts of microbial metabolites as an alternative method to off-line extraction.	Chromatographia, 1984, 19, 246-250.
Moclobemide (monoamine oxidase inhibitor)	Determination of the new monoamine oxidase inhibitor moclobemide and three of its metabolites in biological fluids by HPLC.	J. Chromatogr. Biomed. Appl., 1987, 64 (J. Chromatogr., 420), 110-120.
Mononitroso piperazine	Determination of N-mononitrosopiperazine and N,N-dinitrosopiperazine in human urine, gastric juice and blood.	J. Chromatogr., 1983, 278 (1), 71-80.
Morphine	Combined liquid-solid chromatography and radioimmunoassay for determination of morphine in human fluids.	J. Pharmacol. Methods, 1984, 11 (4), 227-237.

Compound	Title	Journal
Morphine	Concentrations of free and conjugated morphine in blood in twenty cases of heroin-related deaths.	J. Chromatogr. Biomed. Appl., 1987, 66 (J. Chromatogr., 422), 267-273.
Morphine	Determination of morphine in plasma by GC using a macrobore column and thermoionic detection after EXTrelut NT column extraction: application to follow-up morphine treatment in cancer patients.	J. Chromatogr. Biomed. Appl., 1989, 81 (J. Chromatogr., 489), 411-418.
Morphine	Morphine determination by GC-MS in human vitreous humour and comparison with radio-immunoassay.	J. Anal. Toxicol., 1992, 16 (6), 372-374.
Morphine	Comparative study of different extractive procedures to quantify morphine in urine by HPLC - UV.	J. Liq. Chromatogr., 1992, 15 (9), 1533-1541.
Morphine	Comparison of solid-phase extraction and supercritical fluid extraction for the analysis of morphine in whole blood.	J. Anal. Toxicol., 1999, 23(3), 216-218.
Morphine, Amphetamine	Analysis of morphine and amphetamine in meconium with immunoassay and HPLC-diode-array detection.	J. Anal. Toxicol., 1994, 18 (5), 294-295.
Mucoregulatory drug (CO 1408)	HPLC determination of the mucoregulatory drug CO 1408 (-)-2-[6(S)-hydroxy-4(R)-(1-hydroxy-methylethyl)cyclohexenyl]ethanol in rat plasma and urine.	J. Chromatogr., 1990, 507, 227-234.
Nadolol diastereomers	Determination of nadolol diastereomers in dog plasma using chiral derivatization and reversed-phase HPLC with fluorescence detection.	J. Chromatogr. Biomed. Appl., 1994, 661 (2), 281-289.
Nandrolone, Trenbolone	Determination of 19-nortestosterone and trenbolone in animal tissues by HPLC with immunoaffinity clean up.	Food. Addit. Contam., 1998, 15(3), 293-301.
Naphthalene	Naphthalene in biological fluids.	Res. Pract. Forensic Med., 1988, 31, 151-156.
Nicergoline metabolite	Determination of 10-alpha-methoxy-9,10-dihydrolysegol, a nicergoline metabolite, in human urine by HPLC.	Biomed. Chromatogr., 1992, 6 (1), 91.
Nicotine	Improved method for routine determination of nicotine and its main metabolites in biological fluids.	J. Chromatogr., 1982, 232 (2), 295-303.
Nicotine	Microanalysis of nicotine in urine and plasma.	Eisei Kagaku, 1982, 28 (1), 13-17.
Nicotine	Determination of nicotine and four metabolites in the serum of smokers by HPLC with ultra-violet detection.	J. Chromatogr. Biomed. Appl., 1993, 132 (J. Chromatogr., 621), 257-261.
Nicotine	Determination of nicotine and two major metabolites in serum by solid-phase extraction and HPLC and HPLC - particle beam - MS.	J. Chromatogr. Biomed. Appl., 1993, 123 (J. Chromatogr., 612), 209-213.
Nicotine	Rapid solid-phase extraction method for automated GC-MS determination of nicotine in plasma.	J. Chromatogr. Biomed. Appl., 1998, 707(1-2), 312-316.
Nicotine, Cotinine	Simultaneous determination of nicotine and cotinine in urine by GLC.	Eisei Kagaku, 1986, 32 (4), 276-280.
Nimesulide and its metabolites	MS characterization and HPLC determination of the main urinary metabolites of nimesulide in man.	J. Pharm. Biomed. Anal., 1998, 18(1-2), 201-211.
Nimustine hydrochloride	Rapid and simple method for the determination of nimustine hydrochloride in human blood and brain by HPLC.	J. Chromatogr., 1990, 526 (2), 507-514.
Nitenpyram and its metabolites	Determination of the neonicotinoid pesticide nitenpyram and its metabolites in agricultural products.	Shokuhin Eiseigaku Zasshi, 1998, 39(2), 127-134.
Nitrazepam	HPLC determination of nitrazepam and its metabolites in human urine.	J. Chromatogr. Biomed. Appl., 1984, 35 (J. Chromatogr., 310), 213-218.
Nitroglycerin and its dinitrate metabolites	Determination of the two dinitrate meabolites of nitroglycerin in human plasma by capillary GC with electron-capture detection.	J. Chromatogr. Biomed. Appl., 1987, 57 (J. Chromatogr., 413), 101-108.
Nitrophenols	HPLC of nitrophenols determination of products of on-column nitration of phenol.	J. High Resolut. Chromatogr. Chromatogr. Commun., 1987, 10 (11), 624-625.
Nitrosamines	Investigation on occurende of volatile nitrosamines in marinated herrings prepared with and without addition of nitrate.	Z. Lebensm. Unters. Forschung, 1983, 177 (4), 261-263.
N-Nitrosamines, volatile and non-volatile	Quantitative assay of volatile and non-volatile N-nitrosamines by GC with an electrolytic detector. I. Method development and assay of N-nitrosodiethanolamine (NDELA) in creams and lotions.	J. Soc. Cosmet. Chem., 1988, 39 (6), 329-346.
Nitrosamines, volatile	Rapid solid-phase extraction method for the detection of volatile nitrosamines in food.	J. Agric. Food. Chem., 1997, 45(12), 4706-4713.
N-Nitrosodimethylamine	Procedure for GC determination of carcinogenic N-nitrosodimethylamine as a contamination in drugs containing aminophenazone.	Arzneim. Forsch., 1977, 29, 867-869.
Nitrosomethylurea	HPLC determination of N-nitrosomethylurea in biological fluids by reaction with 4-nitrothiophenol.	Food-Chem. Toxicol., 1986, 24 (3), 247-250.
N-Nitrosopiperazine	Determination of N-nitrosopiperazine and N,N'-di-nitrosopiperazine in human urine, gastric juice and blood.	J. Chromatogr. Biomed. Appl., 1983, 29 (J. Chromatogr., 278), 71-80.
Nitrosourea	Qualitative and quantitative analysis of a new sulphur-containing nitrosourea in blood by HPLC.	J. Chromatogr. Biomed. Appl., 1987, 63 (J. Chromatogr., 419), 381-387.
Norethisterone and six metabolites	Simultaneous determination of norethisterone and six metabolites in human plasma by capillary GC with mass-selective detection.	J. Chromatogr. Biomed. Appl., 1995, 674 (2), 155-165.
Norflurazon and oxadixyl residues	Simultaneous extraction of norflurazon and oxadixyl residues from food crops with supercritical carbon dioxide.	Chromatographia, 1996, 42 (3-4), 147-150.
Noscapine	New sensitive method to determine noscapine in serum by reversed-phase LC.	J. Chromatogr. Biomed. Appl., 1985, 38 (J. Chromatogr., 337), 429-433.
Ochratoxin A	Radio-immunoassay of ochratoxin A in barley.	Appl. Environ. Microbiol., 1985, 50 (2), 529-531.
Oestrogens	Determination of free oestrogens in serum using HPLC and electrochemical detection.	Analysis, 1990, 18 (5), 319-322.
Opiates	Procedure for GC determination of opiates in biological samples.	J. Chromatogr., 1981, 206, 109-116.
Organic acids	GC-MS and GC-FT-IR analysis of organic acids: preliminary data.	Biomed. Environ. Mass Spectrom., 1988, 16 (1-2), 285-288.
Organo-chlorine /organo-phosphorus pesticides	Multicolumn solid-phase extraction clean up of organophosphorus and organochlorine pesticide residues in vegetable oils and butterfat.	J. Assoc. Off. Anal. Chem. Int., 1995, 78 (2), 431-437.

Compound	Title	Journal
Organo-chlorine /organo- phosphorous pesticides	Behaviour of 23 persistent organochlorine compounds during sulphuric acid cleanup on a solid-matrix column.	J. Chromatogr., 1990, 513, 33-337.
Organochlorine compounds	Multicolumn solid-phase extraction clean up of organophosphorus and organochlorine pesticide residues in vegetable oils and butterfat.	J. Assoc. Off. Anal. Chem. 1995, 78 (2), 431-437.
Organochlorine compounds	Sandwich-type extraction column with online sulfuric acid treatment for the determination of organochlorine compounds in vegetable oil or oilseeds by GC with electron-capture detection.	J. Chromatogr., 1993, 642 (1-2), 253-262.
Organochlorine pesticides	Solid-matrix partition for separation of organochlorine pesticide residues from fatty materials.	J. Chromatogr., 1991, 552 (1-2), 241-247.
Organochlorine pesticides	Single-step separation of organochlorine pesticide residues from fatty materials by combined use of solid-matrix partition and C18 cartridges.	J. Chromatogr. A., 1997, 765(1), 61-68.
Organophosphate pesticides	On-column partition cleanup of fatty extracts for organophosphate pesticide residue determination.	J. Assoc. Off. Anal. Chem., 1987, 70 (1), 106-108.
Organophosphate pesticides	Single-step multi-cartridge cleanup for organophosphate pesticide residue determination in vegetable oil extracts by GC.	Analyst, 1990, 115 (9), 1167-1169.
Organophosphorous pesticides	Single-step solid-matrix cleanup of vegetable extracts for organophosphorus pesticide residue determination.	J. Chromatogr., 1988, 456 (1), 149-153.
Organophosphorous pesticides	Screening of organophosphorus pesticides using LC-APCI-MS.	J. Chromatogr., 1992, 595, 193-202.
Organophosphorus pesticides	Multiresidue method for determination of organophosphorus pesticides in vegetables.	Food Addit. Contam., 1998, 15(5), 510-517.
Organophosphorus pesticides	Determination of organophosphorus pesticides in foods using an accelerated solvent extraction system.	Analyst, 1997, 122(3), 217-22.
Ornidazole	Simple LC method for the determination of ornidazole and metronidazole in human serum.	J. Chromatogr. Biomed. Appl., 1987, 64 (J. Chromatogr., 420), 228-230.
Ornidazole	Determination of ornidazole and its main metabolites in biological fluids.	J. Chromatogr. Biomed. Appl., 1990, 99 (J. Chromatogr., 534), 233-240.
Oxcarbazepine	Stereoselective bioanalysis of oxcarbazepine and the enantiomers of its metabolites by HPLC.	J. Liq. Chromatogr., 1995, 18 (8), 1533-1541.
Oxidative hair dye compounds	Practical GC-MS analysis of oxidation dye components in hair fibre as a forensic investigative procedure.	J. Forensic. Sci., 1999, 44(2), 292-296.
Oxine-copper	Determination of oxine-copper in organics by GC.	J. Chromatogr., 1989, 474 (2), 381-387.
Oxpentifylline and its metabolite	Determination of 3,7-dimethyl-(5-oxohexyl)xanthine (oxpentifylline) and its 1-(5-hydroxyhexyl)-3,7-dimethylxanthine metabolite in the plasma of patients with multiple diseases using HPLC.	J. Chromatogr. Biomed. Appl., 1984, 30 (J. Chromatogr., 305), 419-427.
Oxprenolol	Determination of oxprenolol in plasma by column LC.	J. Chromatogr. Biomed. Appl., 1988, 70 (J. Chromatogr., 426), 229-233.
Oxprenolol	Identification and quantitative determination of oxprenolol in blood.	J. Planar Chromatogr., 1989, 2 (2), 153-155.
Oxybuprocaine	Oxybuprocaine and five metabolites simultaneously determined in urine by GC and GC-MS after extraction with EXTrelut NT.	Clin. Chem., 1987, 33 (5), 697-700.
Patulin	LC determination of patulin in french apple ciders.	J. Assoc. Off. Anal. Chem., 1996, 79 (5), 1107-1110.
PCDD, PCDF, PCB, PAH	PCDD (polychlorinated dibenzo-p-dioxins), PCDF (polychlorinated dibenzofurans), PCB (polychlorinated biphenyls), PAH (polyaromatic hydrocarbons), cadmium and lead in roadside soil: Relationship between road distance and concentration.	Chemosphere, 1992, 24 (8), 1077-1083.
Pentachloronitro-benzene	LC study of degradation and metabolism of pentachloronitrobenzene by four soil micromycetes.	Chemosphere, 1996, 33 (4), 683-692.
Pesticides	Simplified extraction and cleanup for multi-residue determination of pesticides in lanolin.	J. Assoc. Off. Anal. Chem., 1989, 72 (6), 991-993.
Pesticides	Multiresidue analysis of various kinds of pesticides in cereals by SFE, GC and HPLC.	Shokuhin Eiseigaku Zasshi, 1999, 40(1), 68-74.
Pesticides	Studies on simultaneous determination of fifteen pesticides in cereals by SFE and HPLC.	Shokuhin Eiseigaku Zasshi, 1998, 39(3), 184-191.
PHB esters, Sorbic acid	Simultaneous determination of sorbic acid, dehydroacetic acid, benzoic acid and p-hydroxy-benzoic acid esters by pretreatment with EXTrelut NT column.	Fukushima-ken Eisei Kogai Kenkyusho Nenpo, 1985(3), 100-105.
Phenobarbital	Phenobarbital in hair and drug monitoring.	Forensic Sci. Int., 1995, 70 (1-3), 191-202.
Phenols	Schotten-Baumann benzylation of phenols on an EXTrelut NT column.	J. Chromatogr., 1987, 398, 320-322.
Phenols	Application of liquid-liquid partition chromatography to a student exercise in organic chemistry. IV. Nitration of phenol in an EXTrelut NT column.	Kagaku to Kyoiku, 1987, 35 (3), 251-253.
Phenols, p-Nitrophenols	On-column organic reactions. III. Catalysis by phenol in the nitration of p-nitrophenol with aqueous nitric acid on an EXTrelut NT column.	Chem. Pharm. Bull., 1988, 36 (7), 2661-2664.
Phenytoin	Sensitive method for the determination of phenytoin in plasma, and phenytoin and 5-(4-hydroxyphenyl)-5-phenylhydantoin in urine by HPLC.	J. Pharm. Biomed. Anal., 1992, 10 (10-12), 1001-1006.
Phenytoin, Hydroxyphenytoin	HPLC determination of phenytoin and hydroxyphenytoin in human urine.	J. Chromatogr. Biomed. Appl., 1983, 26 (J. Chromatogr., 275), 97-105.
Phospholipids	Trace analysis of phospholipids by enzymic reactions on a silica-gel plate.	Nippon Kagaku Kaishi, 1990 (8), 896-897.
Phospholipids	Flow-injection analysis for determination of choline-containing phospholipids by luminol chemiluminescence.	Yakugaku Zasshi, 1992, 112 (1), 50-55.
Piperazine	GC method for determination of a piperazine derivative (Trelibet) and its metabolites in human plasma and urine.	J. Chromatogr. Biomed. Appl., 1986, 51 (J. Chromatogr., 378), 155-162.
Piperazine (anti-hypertensive drug)	Determination of the antihypertensive drug 1-[2-ethoxy-2-(3'-pyridyl)ethyl]-4-(2'-methoxy-phenyl) piperazine (IP/66) in rat and human plasma by HPLC and isotope dilution MS.	J. Chromatogr., 1989, 487 (2), 331-340.

Compound	Title	Journal
Piroxicam	HPTLC assay for the routine determination of piroxicam in plasma, urine and tissue.	J. Chromatogr. Biomed. Appl., 1983, 27 (J. Chromatogr., 276), 243-248.
Pirprofen	Determination of the (+)- and (-)-enantiomers of pirprofen in human plasma by HPLC.	J. Chromatogr. Biomed. Appl., 1987, 58 (J. Chromatogr., 414), 131-137.
Pirprofen	HPLC determination of pirprofen and five of its metabolites in human plasma without hydrolysis and in human urine before and after chemical hydrolysis.	J. Chromatogr. Biomed. Appl., 1989, 87 (J. Chromatogr., 495), 195-203.
Polycyclic aromatic hydrocarbons, Heterocyclic aromatic amines	Solid-phase extraction for the selective isolation of polycyclic aromatic hydrocarbons, aza-arenes and heterocyclic aromatic amines in charcoal-grilled meat.	J. Chromatogr. A, 1996, 731 (1-2), 85-94.
Polyethylene glycol 400	HPLC determination of polyethylene glycol 400 in urine: Oligomeric profile in healthy and coeliac disease subjects.	Clin. Chem., 1994, 40 (8), 1571-1574.
Polyphenols	Polyphenolic content in five Tuscany cultivars of <i>Olea europaea</i> L.	J. Agric. Food. Chem., 1999, 47(3), 964-967.
Polysorbates	Determination of polysorbates in foods by colorimetry with confirmation by IR spectrophotometry, TLC and GC.	J. Assoc. Off. Anal. Chem., 1989, 72 (1), 27-29.
Prazepam	Quantitative analysis of prazepam and its metabolites by electron capture GC and selected ion monitoring. Application to dia-placetal passage and fetal hepatic metabolism in early human pregnancy.	J. Chromatogr., 1978, 146 (2), 227-239.
Pregnenolone	Quantitation of pregnenolone and 17-hydroxypreg-nonolone in human serum by automatic HPLC and subsequent radioimmunoassay.	J. Chromatogr., 1981, 224, 229-237.
Preservatives in food	Separation of flavoring agents and preservatives from food by EXTrelut NT.	Nara-ken Eisei Kenkyusho Nenpo, 1980, 15, 86-88.
Procainamide	Simultaneous determination of procainamide and N-acetylprocainamide in serum by GC with nitrogen-selective detection.	J. Chromatogr. Biomed. Appl., 1987, 59 (J. Chromatogr., 415), 143-147.
Progesterone	Procedure for GC determination of progesterone in rat plasma.	J. Chromatogr., 1979, 177, 237-244.
Propylene glycol	Rapid analysis of propylene glycol in fatty foods.	Eisei Kagaku, 34 (2), 1988, 175-178.
Prostaglandin E1	Determination of prostaglandin E1 and its main plasma metabolites 15-ketoprostaglandin E0 and prostaglandin E0 by GC - negative-ion-CI triple-stage.	Biol. Mass Spectrom., 1994, 23 (3), 165-170.
Prothiophos	A case of death from prothiophos poisoning.	Forensic Sci. Int., 1988, 38 (1-2), 13-19.
Proxiphylline	HPLC determination of (R)- and (S)-proxiphylline in human plasma.	J. Chromatogr. Biomed. Appl., 1989, 83 (J. Chromatogr., 491), 355-366.
Psilocin	Quantitation of psilocin in human plasma by HPLC and electrochemical detection: comparison of liquid-liquid extraction with automated on-line solid-phase extraction.	J. Chromatogr. Biomed. Appl., 1998, 709(2), 255-263.
Pyrethrins	Determination of pyrethroid pesticide residues in fatty materials by solid-matrix dispersion partition, followed by mini-column size-exclusion chromatography.	J. Chromatogr., 1999, 833(1), 19-34.
Pyrethrins	Clean-up of aqueous acetone vegetable extracts by solid-matrix partition for pyrethroid residue determination by GC - electron capture detection.	J. Chromatogr., 1997, 765(1), 39-49.
Pyrrolidinone	Determination of pyrrolidinone in plasma by HPLC.	J. Chromatogr. Biomed. Appl., 1985, 45 (J. Chromatogr., 344), 125-136.
Pyrrolizidine alkaloids	Determination of pyrrolizidine alkaloids in honey from selected sites by solid-phase extraction and HPLC-MS.	Food. Addit. Contam., 1997., 14(5), 419-428.
Quinoline	2-amino-3,4-dimethylimidazo-4,5-F-quinoline is not detectable in commercial instant and roasted coffees.	J. Agric. Food Chem., 1991, 39 (12), 2231-2236.
Resorcinol	Determination of free resorcinol in human serum and urine by HPLC with amperometric detection.	Ann. Chim., 1986, 76 , 419-426.
Retention index values	Impact of biological matrix, drug concentration and method of isolation on detectability and variability of retention index values in GC.	J. Anal. Toxicol., 1985, 9 (2), 49-54.
Retinoic acid	Determination of the new aromatic retinoic acid Ro 13-7410 in plasma by two-dimensional GC - negative ion - CIMS with selected-ion monitoring.	Biomed. Environ. Mass Spectrom., 1989, 18 (7), 453-463.
Retinol	Disposable cartridge extraction of retinol and alpha-tocopherol from feedstuffs.	J. Assoc. Off. Anal. Chem., 1985, 68 (6), 1121-1125.
Retinol, alpha-Tocopherol	Disposable cartridge extraction of retinol and alpha-tocopherol from fatty samples.	J. Assoc. Off. Anal. Chem., 1988, 71 (1), 12-15.
Revenast hydrochloride	Determination of revenast hydrochloride in human plasma by GC with nitrogen-selective detection.	J. Chromatogr. Biomed. Appl., 1987, 63 (J. Chromatogr., 419), 351-356.
Rifabutin	Specific HPLC assay for determination of rifabutin plasma concentration following EXTrelut column extraction.	J. Chromatogr. Biomed. Appl., 1999, 728(2), 233-239.
Risperidone, 9-Hydroxyrisperidone	Determination of risperidone and 9-hydroxyrisperidone in human plasma by HPLC with electrochemical detection.	J. Chromatogr. Biomed. Appl., 1993, 125 (J. Chromatogr., 614), 333-339.
Rocuronium and its 17-desacetyl metabolite	Quantification of the neuromuscular blocking agent rocuronium and its putative metabolite 17-desacetylrocuronium in heparinized plasma by capillary GC using a nitrogen sensitive detector.	J. Chromatogr. Biomed. Appl., 1997, 702(1-2), 111-117.
Saccharin	Determination of food preservatives and saccharin by HPLC.	J. Chromatogr., 1979, 173 (2), 343-348.
Sesquiterpene lactones	Sesquiterpene lactones from <i>Arnica chamissonis</i> Less. VI. Identification and quantitative determination by HPLC and GC.	J. Chromatogr., 1987, 410 (2), 329-342.
Sherry-like wine (Vernaccia di Oristano)	Volatile composition of Vernaccia di Oristano sherry-like wine as affected by biological ageing.	J. Sci. Food Agric., 1996, 70 (1), 44-50.
Silybin	LC assay of silybin (silymarin I) in human plasma and urine.	J. Liq. Chromatogr., 1991, 14 (7), 1285-1296.
Steroid hormones	Improved extraction method for plasma steroid hormones.	Clin. Chim. Acta, 1979, 93 (3), 429-431.
Steroid hormones	Simultaneous determination of fifteen steroid hormones from a single serum sample by HPLC and radio-immunoassay.	J. Chromatogr. Biomed. Appl., 1984, 35 (J. Chromatogr., 310), 386-389.
Steroids	Rapid and efficient extraction of steroids from microliter quantities of plasma using EXTrelut NT mini-columns.	Clin. Chim. Acta, 1980, 108 (3), 479-482.
Steroids	Testosterone radioimmunoassay: rapid extraction technique for steroids based on kieselguhr mini-columns.	Steroids, 1980, 36 (2), 149-159.

Compound	Title	Journal
Steroids	A reverse phase HPLC-UV spectrometry method for the analysis of several intrinsic adrenal delta 4-steroid concentration.	Nippon Naibunpi Gakkai Zasshi, 62 (12), 1986, 1326-1335.
Steroids	Semi-automated application of steroids on to TLC sheets.	Anal. Biochem., 1995, 231 (2), 447-449.
Steroids	Improved detection of steroids on NH ₂ layers.	Chromatographia, 1995, 41 (3-4), 175-177.
Steroids	The use of supercritical-fluid extraction for the determination of steroids in animal tissues.	Analyst, 1998, 123(12), 2671-2676.
Steroids, adrenal	Simultaneous quantitative measurement of fourteen adrenal steroids by capillary column GC-MS and its clinical application.	J. Chromatogr., 1986, 374 (1), 5-16.
Sterols	New, rapid method for the separation of labelled sterols synthesized by cell cultures by means of EXTrelut NT.	Fresenius'Z. Anal. Chem., 1985, 322 (5), 512-513.
Strychnine	A new HPLC method for the quantitation of strychnine in urine and tissue extracts.	J. Clin. Chem. Clin. Biochem., 1982, 20 (4), 203-206.
Sulphasalazine	LC method for the simultaneous determination of sulphasalazine and its metabolites in biological fluids.	Chromatographia, 1987, 24, 720-724.
Sulphonamides	LC with amperometric detection of some sulphonamides and their N-4-acetyl-metabolites in serum und urine.	J. Pharm. Biomed. Anal., 1991, 9, 925-928.
Sulthiame	Analysis of sulthiame in serum by narrow-bore HPLC. Comparison of direct sample injection with pre-column switching and EXTrelut NT extraction.	J. Chromatogr., 1991, 553 (1), 7-13.
Synephrine enantiomers	Determination of synephrine enantiomers in food and conjugated synephrine in urine by HPLC.	Anal. Biochem., 1996, 235 (2), 191-194.
Temazepam	Development of a supercritical fluid extraction method for the determination of temazepam in whole blood.	J. Anal. Toxicol., 1997, 21(4), 297-300.
Tenoxicam	Determination of tenoxicam, and the isolation, identification and determination of Ro 17-6661, its major metabolite, in human urine.	J. Chromatogr., 1984, 317, 483-492.
Terpenes	Analysis of terpenes from Ginkgo Biloba L. extracts by RP-HPLC.	Chromatographia, 1990, 29, 251.
Tertatolol	Determination of tertatolol enantiomers in biological fluids by HPLC.	J. Chromatogr. Biomed. Appl., 1991, 110 (J. Chromatogr., 572), 203-210.
Testosterone	Measurement of plasma testosterone by GC - negative ion - CIMS using pentafluoropropionic derivatives.	J. Chromatogr. Biomed. Appl., 1995, 663 (2), 187-192.
Testosterone	Determination of testosterone in human hair by GC - selectedion monitoring - MS.	Analyst, 1998, 123(12), 2661-2663.
Testosterone, Androstadienone	Simultaneous determination of testosterone and androstadienone (sex attractant) in human plasma by GC-MS with high-resolution selected-ion monitoring.	J. Chromatogr. Biomed. Appl., 1991, 103 (J. Chromatogr., 565), 35-44.
Tetraorganolead compounds	Metabolism of tetraorganolead compounds by rat-liver microsomal monooxygenase. I. Analytical methods for quantitative determination of dealkylation products.	Xenobiotica, 1983, 13 (10), 577-581.
Theophylline, Theobromine, Caffeine	Highly sensitive and rapid determination of theophylline, theobromine and caffeine in human plasma and urine by gradient capillary HPLC - frit - FABMS.	J. Chromatogr. Biomed. Appl., 1995, 667 (2), 241-246.
Thiabendazole	Capillary GC determination of thiabendazole in citrus and apple juices.	J. Assoc. Off. Anal. Chem., 1994, 77(5), 1293-1296.
Thiamphenicol	Procedure for LC determination of thiamphenicol in bovine serum and milk.	J. Liq. Chromatogr., 1994, 17 (12), 2625-2632.
Thiopental	Toxicological study on intravenous thiopental anesthesia-interrelation between rate of injection and distribution of thiopental.	Forensic Sci. Int., 1987, 33 (3), 169-175.
Ticlopidine	Quantitative HPLC, GC, and GC-MS analysis of ticlopidine in baboon plasma after solid-phase extraction.	J. Pharm. Sci., 1991, 80 (11), 1092-1095.
Timelotem dihydrogen maleate	Determination of timelotem dihydrogen maleate and its main metabolite N-desmethyltimelotem in plasma by capillary GC.	J. Chromatogr. Biomed. Appl., 1987, 63 (J. Chromatogr., 419), 357-362.
Tiropamide	Identification of metabolites of tiropamide in human urine.	Biomed. Environ. Mass Spectrom., 1988, 15 (4), 205-209.
Toluene-2,4-diamine	GC determination of 2,4-toluenediamine (toluene-2,4-diamine) in urine.	Ann. Chim., 1992, 82 (3-4), 179-183.
Trace aroma and flavour components	Solid-phase extraction and determination of trace aroma and flavour components in cider by GC-MS.	Chromatographia, 1996, 42 (1-2), 101-105.
Trace organic impurities	Isolation, characterization and determination of trace organic impurities in FD and C Red No. 40 [C.I. Food Red 17].	J. Chromatogr., 1989, 467, 167-176.
Tranquilizers	GC and GC/MS analysis of tranquilizers in meat.	Fresenius' Z. Anal. Chem., 1982, 311, 398-399.
Triclosan	Selected-ion-monitoring assay for triclosan in medical waste water.	Biomed. Environ. Mass Spectrom., 1988, 16, 255-257.
Triethylene thiophosphoramidate	HPLC of the anti-tumour agent triethylene thiophosphoramidate and its metabolite triethylene phosphoramidate with sodium sulphide, taurine and phthalaldehyde as pre-column fluorescent derivatization reagents.	J. Chromatogr., 1988, 458, 295-301.
Triethylene thio-phosphoramidate	Spectrofluorimetric determination of triethylene thiophosphoramidate in blood.	Anal. Chim. Acta, 1987, 201, 77-82.
Vitamin D ₃	A sensitive and simplified radioimmunoassay for 1alpha,25-dihydroxyvitamin D ₃ .	Clin. Chim. Acta, 1984, 142 (3), 325-338.
Vitamin D ₃	Rapid method for measuring physiological concentrations of 25-hydroxyvitamin D levels in blood serum.	Clin. Chim. Acta, 1985, 153 (2), 85-92.
Yutac	HPLC method for quantitative determination of Yutac, a new antiarrhythmic agent, in dog plasma.	J. Chromatogr. Biomed. Appl., 1985, 39 (J. Chromatogr., 338), 428-432.
Zindoxifene	Chromatographic analysis of the new antiestrogen zindoxifene and its metabolites in biological material.	J. Chromatogr., 1987, 414, 235-241.

エクストレルート NT 製品ラインアップ

プレパックカラム

製品名	サンプル容量 (mL)	カラム材質	カラム数 (一箱中)	注文番号
エクストレルート NT1	0.1-1	ガラス	100	1.15094.0001
エクストレルート NT3	1-3	ガラス	50	1.15095.0001
エクストレルート NT20	<20	プラスチック	25	1.15096.0001

詰め替え用充填剤

製品名	包装単位	注文番号
エクストレルート NT20 用 詰め替えパック	50 パック	1.15093.0001
エクストレルート NT 充填剤 (詰め替え用)	1kg	1.15092.0001

アクセサリー

製品名	サイズ	包装単位	注文番号
エクストレルート NT1/NT3 用 カニューレ	0.6 × 30 mm	100 本	1.15373.0001
エクストレルート NT1/NT3 用 コレクションチューブ	15 ml タイプ	30 本	1.15622.0001
交換フィルター NT1 用	直径 10 mm	100 枚	1.14236.0001
交換フィルター NT3 用	直径 15 mm	100 枚	1.14237.0001
交換フィルター NT20 用	直径 24 mm	50 枚	1.14567.0001

高分解能 HPLC カラム

ピューロスフェア STAR

Purospher® STAR

- 高カラム効率
- 優れた選択性
- 再現性
- 高い耐久性

ピューロスフェア STAR はメルクが現在提供している最新の HPLC 担体です。メルク製品ラインアップの中で最も高い理論段数を実現しつつ、広範囲の pH で利用可能です。担体にはきわめて純度の高い球状シリカゲルを使用しています。総金属不純物含量は 5ppm 以下に抑えられており、酸性・塩基性、キレート化合物を問わず非常に美しいクロマトグラムが得られます。

● ピューロスフェア STAR の主なスペック

金属含有量：Na, Ca, Mg, Al: 1ppm 以下、Fe: 3ppm 以下
 細孔径：120Å 比表面積：330m²/g
 使用可能 pH 範囲：1.5 - 10.5(RP-8e・RP-18e)
 2.0 - 7.5(Si・NH₂)

pH 安定性

Stability test at pH 10.5

Stability test at pH 1.5

● 製品ラインアップ

カラム内径 (mm) \ カラム長 (mm)	0.5	1	2	3	4	4.6	10	21.2	25
30			RP-18e (3 μm)		RP-18e (3 μm)		RP-8e/18e (5 μm)	RP-8e/18e (5 μm)	RP-8e/18e (5 μm)
50							RP-8e/18e (5 μm)	RP-8e/18e (5 μm)	RP-8e/18e (5 μm)
55			RP-18e (3 μm)		RP-18e (3 μm)				
75					RP-18e (3 μm)				
100							RP-8e/18e (5 μm)	RP-8e/18e (5 μm)	RP-8e/18e (5 μm)
125			RP-8e/18e (5 μm)	RP-8e/18e (5 μm)	RP-8e/18e (5 μm)				
150	RP-18e (3 μm)	RP-18e (3/5 μm)				RP-8e/18e/NH ₂ /Si (5 μm)			
250			RP-8e/18e (5 μm)	RP-8e/18e (5 μm)	RP-8e/18e/NH ₂ /Si (5 μm)	RP-8e/18e/NH ₂ /Si (5 μm)	RP-8e/18e (5 μm)		

■ 汎用分析カラム ■ セミ分取カラム ■ ハイスルーブットカラム ■ セミマイクロカラム

メルク株式会社

パフォーマンス・ライフサイエンス化学品事業部

〒153-8927

東京都目黒区下目黒 1-8-1 アルコタワー5F

Tel:0120-189-390 / Fax:0120-189-350

E-mail:service@merck.co.jp

<http://www.merck.co.jp>

平成17年11月21日発行 REA51-0511-5000